

REVISTA NICARAGUENSE DE BIODIVERSIDAD

N° 119

Enero 2025

A review the diet of *Leptodeira rhombifera* (Colubridae) and new case to predation attempt of *Rhinella horribilis* (Bufonidae) in Nicaragua.

Lester Fonseca-González, Ariel Salinas &
Ronier Francisco Rugama Montoya.



PUBLICACIÓN DEL MUSEO ENTOMOLÓGICO
LEÓN - - - NICARAGUA

La Revista Nicaragüense de Biodiversidad (ISSN 2413-337X) es una publicación que pretende apoyar a la divulgación de los trabajos realizados en Nicaragua en este tema. Todos los artículos que en ella se publican son sometidos a un sistema de doble arbitraje por especialistas en el tema.

The Revista Nicaragüense de Biodiversidad (ISSN 2413-337X) is a journal created to help a better divulgation of the research in this field in Nicaragua. Two independent specialists referee all published papers.

Consejo Editorial

Jean Michel Maes
Editor General
Museo Entomológico
Nicaragua

Milton Salazar
Herpetonica, Nicaragua
Editor para Herpetología.
herpingnicaragua@gmail.com

Eric P. van den Berghe
ZAMORANO, Honduras
Editor para Peces.

Liliana Chavarría
ALAS, El Jaguar
Editor para Aves.

José G. Martínez-Fonseca
Nicaragua
Editor para Mamíferos.

Oliver Komar
ZAMORANO, Honduras
Editor para Ecología.

Estela Yamileth Aguilar
Álvarez
ZAMORANO, Honduras
Editor para Biotecnología.

Indiana Coronado
Missouri Botanical Garden/
Herbario HULE-UNAN León
Editor para Botánica.

URL DE LA REVISTA: <http://www.bio-nica.info/revistanicarague/index.html>



Esta obra está bajo una Licencia Creative Commons Atribución - No Comercial - Sin Obra Derivada 4.0 Internacional

Foto de portada: *Leptodeira rhombifera* (Foto © Lester Fonseca).

A review the diet of *Leptodeira rhombifera* (colubridae) and new case to predation attempt of *Rhinella horribilis* (Bufonidae) in Nicaragua.

Lester Fonseca-González¹, Ariel Salinas² & Ronier Francisco Rugama Montoya³.

Resumen

Revisión de la dieta de *Leptodeira rhombifera* (Colubridae) y nuevo caso de intento de depredación de *Rhinella horribilis* (Bufonidae) en Nicaragua.

Las interacciones presa-depredador son importantes para comprender la dinámica de especies, por lo cual aportamos datos relevantes de la dieta de *Leptodeira rhombifera* a través de la búsqueda exhaustiva de publicaciones concerniente a sus presas, conjuntamente reportamos por primera vez el intento de depredación de *Rhinella horribilis* por *L. rhombifera* en el pacífico de Nicaragua. Corroboramos que la dieta de *L. rhombifera* está dirigida hacia anfibios; sin embargo, los reptiles y peces forma parte sustancial de sus hábitos alimenticios. Por último, destacamos que *R. horribilis* podría representar una alternativa en los hábitos alimentarios del género *Leptodeira* y probablemente de otras serpientes terrestres.

Palabras claves: Anfibios, Dieta, Dinámica ecológica, Presa, Serpientes.

DOI: 10.5281/zenodo.14611330

¹ Licenciado en gerencia ambiental y de los Recursos naturales, Universidad Nacional Autónoma de Nicaragua (UNAN - Managua). ORCID: 0000-0002-9886-4813

² Reserva Natura, Responsable de Biodiversidad.

³ Universidad Nacional Agraria (UNA) ORCID: 0009-0002-4139-0060

Abstract

Predator - prey interactions are important for understanding species dynamics; therefore, we provide relevant data on the diet of *Leptodeira rhombifera* through an exhaustive literature review of its diet. We also report for the first time the predation attempt of *Rhinella horribilis* by *L. rhombifera* in the Pacific region of Nicaragua. We corroborate that the diet of *L. rhombifera* primarily targets amphibians; however, reptiles and fish also form a substantial part of its feeding habits. Finally, we emphasize that *R. horribilis* could represent an alternative in the feeding habits of the genus *Leptodeira* and probably of other terrestrial snakes.

Keywords: Amphibians, Diet, Ecological dynamics, Prey, Snakes.

Introduction

Tropical habitats are characterized by high biodiversity and complex biotic interactions such as predation (Freestone *et al.*, 2011; Brown, 2014). Predation is an integral part of the ecological dynamics of living organisms and is based on the effort made by an animal to find and feed on another (Curio, 1976), it is fundamental for better understanding the ecology of living organisms (Rosenberg and Cooper, 1990), as documenting predation interactions provides valuable information to increase the understanding of the dynamics of structure and function within communities and ecosystems (Rosenberg & Cooper, 1990).

In this sense, anurans are a crucial component in the trophic dynamics of communities and ecosystems (Whiles *et al.*, 2006); they are secondary consumers in the food chain, as they consume insects and other invertebrates, while also serving as prey for other species.

Leptodeira is a genus of snakes known for basing its diet mainly on amphibians and small lizards, in *L. rhombifera*, it probably consists mainly of adult or juvenile anurans (Espinoza, 2021; Duellman, 1958; Solórzano, 2004). However, the species has been documented feeding on other groups such as fish (Solórzano, 2022), birds and small mammals (unspecified), (Köhler 2008; Solórzano, 2004). Additionally, there are records mentioning other important aspects of the species diet such as scavenging (Fuentes & Quiroz-Espinoza, 2024; Knight, 2016) and ophiophagy (Köhler 2008; Solórzano, 2004).

L. rhombifera, has an established distribution from southern Mexico to Panama and is found in dry to humid lowland tropical environments (Barrio-Amorós, 2019). In Nicaragua, it is present throughout almost the entire national territory, associated wetlands (HerpetoNica, 2015) and is primarily terrestrial and nocturnal, according to Leenders, (2019) it is more common in seasonally dry areas.

The species possesses a certain amount of venom and enlarged, striated posterior maxillary teeth that they use to immobilize its prey before consuming them (Leenders, 2019).

In another order, Cane toads are the most widely distributed and abundant new world amphibians around the world (Lever 2001), within this group we find *Rhinella horribilis*, which is naturally distributed from southern Texas to the central Amazon. Particularly, this species is one of the most cosmopolitan anurans and inhabits diverse types of ecosystems (Acevedo *et al.*, 2016; Bonett *et al.*, 2017). In Nicaragua, it is present throughout the country, in urban areas, livestock areas and altered natural habitats (HerpetoNica, 2015). It is terrestrial and nocturnal (Savage, 2002) although arboreal behavior has been reported (Berra, 2020).

This toad has a generalist diet, feeding on a wide variety of invertebrates, small vertebrates and in some cases, plants (Botero-Trujillo 2006; Escudero & Ortega 2009; Sampedro-Marín *et al.*, 2011), essentially consuming different species that it can capture in its mouth.

In this note, we describe a case attempted predations of *R. horribilis* by *L. rhombifera* in the pacific region of Nicaragua, and discuss relevant aspects of the diet of *L. rhombifera*.

Materials and Methods

The event of predation occurred in the central Pacific region of Nicaragua, department of Managua, in “Natura” Reserve, located at the following coordinates: N. 11.86756°, W. 086.51387°. This zone corresponds to Central American dry forests and secondary regenerations was present locally.

Information gathering

To document the in-situ observation and identification of both species involved, the organisms were photographed using a Sony Cyber-shot H-400 and Nikon 3500 camera, without interrupting the event from a distance of one meter.

An exhaustive search for publications concerning the prey of *L. rhombifera* was conducted using the Google Scholar search engine and Scopus. We also reviewed the metadata of 15 specialized herpetology journals, as well as the repository of CSUCA (Consejo superior de Universidades Centroamericanas) and other academic institutions, primarily in North America. During the search we delimited the cases explicitly documented for *L. rhombifera* and employed field operators (Gómez, 2017) with the key terms: Alimentación / Dieta / Depredación / Nota / *Leptodeira rhombifera* / *Rhinella horribilis* / Registro / Nota, as well as their English equivalents.

The compiled information was carefully reviewed and selected from its source and was organized chronologically and by taxonomic group.



Figure 1. Location of predation case.

Results and Discussion

On August 20, 2024, during a night walk in the vicinity of the “Los Lagartos” reservoir, a juvenile individual of *L. rhombifera* was observed on one side of the “Guardabarranco” trail, feeding on a specimen of juvenile *R. horribilis*. The event took place at 18:05 hours, we proceeded to photograph the predation for no more than four minutes during which the snake attempted to swallow the frog (figure 2). At the time of the observations, the snake was holding its prey by the right flank, the prey showed no signs of resistance.

Later, we moved to other locations within the reserve, when returning to side, we observed that the snake abandoned its prey, the anuran exhibited lacerations on the shoulders as a result of the snake's subjection.




Figure 2. Event of predation of *R. horribilis* by *L. rhombifera*. (photos © Lester Fonseca and Ariel Salinas).

We cannot assume that the colubrid did not consume the prey due to our presence or because it could not swallow it, as such cases have been previously recorded in snakes of the Colubridae family, which do not complete their feeding for different reasons (Heinen & Hammond, 1997; Engeman & Engeman, 2015; Teles *et al.*, 2018; Hernández & Guevara, 2022). This case may be related to the relative sizes of the predator and prey, or to secretions on the skin of the frogs, as well as to physical resistance on the part of the prey (Costa & Trevelin, 2020).

Both the prey and the predator (approximately 29 cm) were juveniles, but the result was the lifeless body of the prey, with torn shoulders and symptoms of having been poisoned (figure 3).



Figure 3. Post mortem state of *R. horribilis* (photos © Ariel Salinas and Lester Fonseca).

Rhinella horribilis has been documented as prey for other species of the genus *Leptodeira*, such as *L. maculata* and *L. splendida* (García-Mata *et al.*, 2017; Gámez-Duarte *et al.*, 2024). Also, other reptiles such *Ctenosaura similis* (Pasachnik & Corneil, 2011), even data provided by Pasachnik & Corneil, (2011) and Okada-Aguiar & Costa-Campos (2018), specify that *R. horribilis* is a feeding alternative even in the post-mortem state.

It is worth mentioning that according to HerpetoNica (2015), the Bufonidae family is considered the most common family in the Pacific region of Nicaragua and is present throughout the country. Within this group, it is important to note that *R. horribilis* is the most frequently observed species. This statement is consistent with the relative abundance of *R. horribilis* in the

“Nature” Reserve, as it tends to be the most commonly observed amphibian. Therefore, this species could represent an alternative resource in the diet of terrestrial snakes due to its common character.

The diet of *L. rhombifera* is well documented, however, records documenting its prey at the species level are rare. For example, (Savage, 2002) mentions that it consumes aquatic prey such as crabs and fish, other authors include lizards, amphibians, tree frog eggs, and small mammals (Duellman, 1958, Solórzano, 2004; Köhler, 2008), although without precision about species.

In this regard, we compile the cases that specifically document the species that are part of the diet of *L. rhombifera*, according our documentary review, the diet of *L. rhombifera* is more focused on amphibians.

The feeding habits by *L. rhombifera* are closely related to its natural history; being a nocturnal and terrestrial species, it relies on species that can be easily found on the ground or in not so high places. On the other hand, these characteristics may explain why there are almost no literary resources documenting the predation of birds and mammals. In addition, it is likely that it resorts to scavenging more often than believed. The consumption of dead prey in *L. rhombifera* is important, as it alludes to an ability of the species to select a broader diet when anurans are less abundant (Céspedes, & Abarca, 2014).


Of the 20 cases collected of predation by *L. rhombifera*, 60% correspond to amphibians, while 20% correspond to reptiles and 20% to fish and we include in their diet small mammals, which have been mentioned without specifying the species. It should be noted that five cases come from experiments, some of which were conducted outside the natural habitat of the species, and four species were documented as cases of scavenging. (figure 4).

Table 1. Compilation the diet (Preys) of *Leptodeira rhombifera*.

Documented prey of <i>Leptodeira rhombifera</i>				
Taxonomic group	Prey	Documented attack	Country	Sources
Amphibians	<i>Engystomops pustulosus</i>	Yes (Individuo)	Barra Colorado, Panama	Ryan (1985)
Amphibians	<i>Engystomops pustulosus</i>	Yes (Individuo)	Gamboa, Panama.	Dougherty & Lisondro (2023)
Amphibians	<i>Boana rosenbergi</i>	Yes (Individuo)	Panama	Kluge (1981)
Amphibians	<i>Boana rosenbergi</i>	Yes (Individuo)	Gamboa, Panama.	Dougherty & Lisondro (2023)
Amphibians	<i>Engystomops pustulosus</i>	Yes (Individuo)	Gamboa, Panama.	Dougherty & Lisondro (2023)
Amphibians	<i>Incilius coccifer</i>	Not (Individuo)	Guanacaste, Costa Rica	Área de Conservación de Guanacaste
Amphibians	<i>Lithobates vallanti</i>	Yes (Scavenging)	Guanacaste, Costa Rica	Mora-Benavides (1999)
Amphibians	<i>Rhinophryne dorsalis</i>	Yes (Individuo)	Guanacaste, Costa Rica	Céspedes et al. (2018)
Amphibians	<i>Anaxyrus americanus*1</i>	Yes (Experiment)	Arrived in New York, EEUU	Skeha (1959)
Amphibians	<i>Lithobates pipiens*2</i>	Yes (Experiment)	Arrived in New York, EEUU	Skeha (1959)
Amphibians	<i>Pseudacris regilla</i>	Yes (Experiment)	EEUU	Busack (2018)
Amphibians	<i>Hylidae</i>	Yes (Scavenging)	Los Santos, Panama	Knight (2016)
Reptiles	<i>Leptodactylus labialis</i>	Yes (Scavenging)	Panama	Fuentes & Quiroz-Espinoza (2024)
Reptiles	<i>Lampropeltis triangulum</i>	Yes (Experiment)	Arrived in New York, EEUU	Skeha (1959)

Reptiles	<i>Anolis spp</i>	Yes (Experiment)	Arrived in New York, EEUU	Skeha (1959)
Reptiles	<i>Iguana rhinolopha</i>	Yes (Scavenging)	Panama	Fuentes & Quiroz- Espinoza (2024) Rojas- Carranza & Anderson (2023)
Fish	<i>Rhamdia guatemalensis</i>	Yes (2 Individuo)	Guanacaste, Costa Rica.	Céspedes & Abarca (2014) Knight & Shervette (2022) Solis & Guerrero (2016)
Fish	<i>Rhamndia spp</i>	Yes (Individuo)	Alajuela, Costa Rica	
Fish	<i>Characidae</i>	Yes (Individuo)	Los santos, Panama	
Fish	<i>Rhamdia laticauda</i>	Yes (Individuo)	Francisco Morazán, Honduras	

Note: * indicates update in the scientific name of some documented species:
*Bufo americanus**¹, *Rana pipiens**².



Our prey collection for *L. rhombifera* suggests a higher degree of plasticity than previously reported (Espinoza, 2021; Duellman, 1958; Solórzano, 2004; Dougherty & Lisondro, 2023).

Conclusion

We confirm that anurans are the main food item of *L. rhombifera*, but this review suggests that fish and reptiles could be an essential part of its diet. Additionally, it is important to mention that the species presents some behaviors that are little known or sometimes overlooked, such as ophiophagy (Köhler 2008; Solórzano, 2004) and scavenging (Fuentes & Quiroz-Espinoza, 2024; Knight, 2016).

Finally, we emphasize that due to the frequency and abundance of *R. horribilis*, this species could represent an alternative in the feeding habits of many predators, including snakes of the genus *Leptodeira* (García-Mata *et al.*, 2017; Gámez-Duarte *et al.*, 2024) and other terrestrial snakes.

Literatura Citada

- Acevedo A.A., Lampo M. & Cipriani R. (2016) The cane or marine toad, *Rhinella marina* (Anura, Bufonidae): Two genetically and morphologically distinct species. Zootaxa. 4103(6): 574-586. <https://doi.org/10.11646/zootaxa.4103.6.7>
- Área de Conservación Guanacaste. Fuente de Vida y Desarrollo (2024) *Leptodeira rhombifera* intenta comer un *Incilius coccifer*. Jairo Moya Vargas. <https://www.acguanacaste.ac.cr/biodiversidadacg/observaciones-de-biodiversidad/4669-leptodeira-rhombifera-intenta-comer-un-incilius-coccifer>
- Barrio-Amorós C.L. (2019) On the taxonomy of snakes in the genus *Leptodeira*, with an emphasis on Costa Rican species. Reptiles & Amphibians. 26: 1-15.
- Berra D. (2020) Comportamiento arbóreo del Sapo de Caña Mesoamericano (*Rhinella horribilis*) en Coquimatlán, Colima, México. Revista Latinoamericana de Herpetología. 3(1): 90-92. <https://doi.org/10.22201/fc.25942158e.2020.1.123>
- Bonett R.M., Boundy J., Burbrink F.T., Crother B.I., Queiroz K., Frost D.R. & Krysko K.L. (2017) Scientific and Standard English Names of Amphibians and Reptiles North of Mexico, with Comments Regarding Confidence in Our Understanding. Referencia ?
- Brown J.H. (2014) Why are there so many species in the tropics? Journal of Biogeography. 41(1):8-22 <https://doi.org/10.1111/jbi.12228>

- Botero-Trujillo R.** (2006) Anuran predators of scorpions: *Bufo marinus* (Linnaeus, 1758) (Anura: Bufonidae), first known natural enemy of *Tityus nematochirus* Mello-Leitão, 1940 (Scorpiones: Buthidae). Revista Ibérica de Aracnología. 13:199-202.
- Busack S.** (2018) *Leptodeira rhombifera* (common cat-eyed snake). Longevity and aging in captivity. Herpetological Review. 49(1): 55.
- Céspedes, J. & Abarca J.** (2014) Nature Notes. *Leptodeira rhombifera*. Diet. Mesoamerican Herpetology. 1(2) 288-289.
- Céspedes J., Astorga J.D., Sánchez J. & Obando J.** (2018) Predation of *Rhinophryne dorsalis* (Duméril and Bibron, 1841) (Anura: Rhinophryidae) by *Leptodeira rhombifera* (Günther, 1872) (Serpentes: Dipsadidae), in Guanacaste, Costa Rica. Herpetology Notes. 11: 959-960.
- Costa W.P. & Trevelin C.C.** (2020) Unsuccessful predation attempts by snakes on anuran amphibians: How successful are snakes? Herpetology Notes. 13:649-660.
- Curio E.** (1976) The Ethology of Predation. New York: Springer Science & Business Media.
- Dougherty R.P. & Lisondro A.K.** (2023) Predation of the anurans *Agalychnis callidryas*, *Boana rosenbergi*, and *Engystomops pustulosus* by the snakes *Leptodeira ornata* and *Leptodeira rhombifera* in an artificial pond. Herpetology Notes. 16: 507-516.
- Duellman W.E.** (1958) A monographic study of the colubrid snake genus *Leptodeira*. Bulletin of the American Museum of Natural History. 114: 1-183.
- Engeman R.M. & Engeman C.** (2015) *Leptodeira septentrionalis* (northern cat-eyed snake). Diet and predation. Herpetological Review. 46(1): 104-105.
- Escudero M.B. & Ortega A.M.J.** (2009) Dieta Entomofágica en Una Población de *Rinella Marina* en La Selva Pluvial Central, Chocó, Colombia. Revista Institucional Universidad Tecnológica Del Chocó Investigación Biodiversidad Y Desarrollo. 28(2).
- Espinosa A.** (2021) Dieta de *Leptodeira* (Colubridae: Serpentes) (Fitzinger 1843) en Ecuador y notas ecológicas de una población de *L. septentrionalis larcorum* (Kennicott 1859) en Zapotillo-Loja, Ecuador. Quito: UCE. Unpublished thesis, Universidad Central del Ecuador, Quito, Ecuador.

Freestone A.L., Osman R.W., Ruiz G.M. & Torchin M.E. (2011) Stronger predation in the tropics shapes species richness patterns in marine communities. *Ecology*. 92(4):983-993 <https://doi.org/10.1890/09-2379.1>

Fuentes R. & Quiroz-Espinoza M. (2024) Necrophagy in *Leptodeira rhombifera* (Squamata: Serpentes) in Panama. *Revista Científica Vida Natural*. 2(1). <https://revistas.unachi.ac.pa/index.php/vidanatural/article/view/764>

Gomez-Duarte E.A., Jacobo J.D., Manríquez J.M., Castro H.A. & Serrano, J.M. (2024) Necrofagia en dos especies de serpientes del género leptodeira (squamata: d ipsadidae). *revista latinoamericana de herpetología*. 7(2): e942 (123 - 128).

García-Mata E.S., Cruz-Saenz D., Carlos-Gomez J.A., Navarro-Velazquez D.L. & Wilson L.D. (2017) Notes on the Herpetofauna of Western Mexico 17: Predation on *Rhinella horribilis* (Linnaeus, 1758) by two species, *Leptodeira maculata* (Hallowell, 1861) and *Caracara cheriway* (Jacquin, 1784), in the municipality of Cuauhtemoc, Colima, Mexico. *Bulletin of the Chicago Herpetological Society*. 52(8):139-145.

Gómez J.A. (2017) Técnicas para el proceso de búsqueda, acceso y selección de información digital: los operadores. *Publicaciones Didácticas*. 87: 393-529.

Heinen J.T., & Hammond G. (1997) Antipredator Behaviors of Newly Metamorphosed Green Frogs (*Rana clamitans*) and Leopard Frogs (*R. pipiens*) in Encounters with Eastern Garter Snakes (*Thamnophis s. sirtalis*). *The American Midland Naturalist*. 137(1): 136-144

Hernández J.J. & Guevara N. (2022) evento de depredación en *Trachycephalus vermiculatus* (anura: hylidae) por *leptophis ahaetulla* (squamata: colubridae) en la provincia de coclé, república de panamá. *revista latinoamericana de herpetología*. 5(1): 56-59.

HerpetoNica (2015) Guía Ilustrada de los Anfibios y Reptiles de Nicaragua. MARENA - HerpetoNica. Managua, Nicaragua.

Knight J.L. (2016) Natural history notes. *Leptodeira rhombifera* (Common Cat-eyed Snake). Neonate diet/scavenging. *Herpetological Review*. 47(2): 313-314.

Knight J.L. & Shervette V.R. (2022) *Leptodeira rhombifera* (common cat-eyed snake; ojo de gato). diet. *Herpetological Review*. 53(3): 513.

- Kluge A.** (1981) The life history, social organization, and parental behavior of *Hyla rosenbergi* Boulenger, a nestbuilding gladiator frog. Miscellaneous Publications Museum of Zoology, University Michigan. 41(160):1-170.
- Köhler G.** (2008) Reptiles of Central America. Second Edition. Offenbach, Germany, Herpeton.
- Leenders T.** (2019) Reptiles of Costa Rica. A Field Guide. A Zona Tropical Publication. Comstock Publishing Associates, Cornell University Press, Ithaca, New York, USA.
- Lever C.** (2001) The cane toad: the history and ecology of a successful colonist. Westbury Academic & Scientific Pub.
- Mora-Benavides J.M.** (1999) Natural Historical Notes: Serpentes. Herpetological Review. 30(2) 102.
- Okada-Aguiar K. & Costa-Campos C.** (2018) *Rhinella marina* (cane toad). scavenged by Caracara. Herpetological Review. 49(2):310-311.
- Pasachnik S.A. & Corneil J.P.** (2011) *Ctenosaura similis* (Black Spinytailed Iguana). Diet. Herpetological Review. 42 (4): 601-602.
- Rojas-Carranza A.H. & Anderson N.** (2023) Predation by the Common Cat-eyed Snake, *Leptodeira rhombifera* Günther, 1872, on the Pale Catfish in Costa Rica. Herpetology Notes.16: 561-563.
- Rosenberg K.V. & Cooper R.J.** (1990) Approaches to avian diet analysis. Studies in Avian Biology. 13: 80-90. <https://sora.unm.edu/node/139273>
- Ryan M.J.** (1985) The Tungara Frog: A Study in Sexual Selection and Communication. University of Chicago Press, Chicago, Illinois, USA.
- Sampedro-Marín, A.C., Angulo Y.Y., Arrieta F.I. & Dominguez D.M.** (2011) Alimentación de *Bufo marinus* (Linnaeus, 1758) (Bufonidae: Anura), en una localidad de Sucre, Colombia. Caldasia. 33(2):495-505.
- Savage J.M.** (2002) The amphibians and reptiles of Costa Rica: A herpetofauna between two continents, between two seas. University of Chicago Press, Chicago, USA, 934.
- Skehan P.** (1959) Ophiophagy in *Leptodeira*. Herpetologica. 15(3):160
- Solís J.M. & Guerrero F.** (2016) Natural history notes. *Leptodeira rhombifera*. Diet. Herpetological Review. 47(2): 313.
- Solórzano A.** (2004) Serpientes de Costa Rica: Distribución, Taxonomía e Historia Natural. Instituto Nacional de Biodiversidad, San José, Costa Rica.

Solórzano A. (2022) Snakes of Costa Rica: Distribution, Taxonomy, and Natural history. Second Edition. San José, Costa Rica, Editorial.

Teles A., De Sena A. & Ribeiro M.V. (2018) Predation attempt of *Xenopholis undulatus* (Serpentes, Dipsadidae) on *Physalaemus cuvieri* (Amphibia, Leptodactylidae). Herpetology Notes. 11: 829-830.

Whiles M.R., Lips K.R., Pringle C. M., Kilham S.S., Bixby R.B., Brenes R., Connelly S., Colon-Gaud J.C., Hunte-Brown M., Huryn A.D., Montgomery C. & Peterson S. (2006) The effects of amphibian population declines on the structure and function of Neotropical stream ecosystems. 4(1): 27-34
[https://doi.org/10.1890/1540-9295\(2006\)004\[0027:TEOAPD\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)004[0027:TEOAPD]2.0.CO;2)

BORRADOR

La Revista Nicaragüense de Biodiversidad (ISSN 2413-337X) es una publicación de la Asociación Nicaragüense de Entomología, aperiódica, con numeración consecutiva. Publica trabajos de investigación originales e inéditos, síntesis o ensayos, notas científicas y revisiones de libros que traten sobre cualquier aspecto de la Biodiversidad de Nicaragua, aunque también se aceptan trabajos de otras partes del mundo. No tiene límites de extensión de páginas y puede incluir cuantas ilustraciones sean necesarias para el entendimiento más fácil del trabajo.

The Revista Nicaragüense de Biodiversidad (ISSN 2413-337X) is a journal of the Nicaraguan Entomology Society (Entomology Museum), published in consecutive numeration, but not periodical. RNB publishes original research, monographs, and taxonomic revisions, of any length. RNB publishes original scientific research, review articles, brief communications, and book reviews on all matters of Biodiversity in Nicaragua, but research from other countries are also considered. Color illustrations are welcome as a better way to understand the publication.

BOLETA DE INFORMACIÓN
Todo manuscrito para RNB debe enviarse en versión electrónica a:
(Manuscripts must be submitted in electronic version to RNB editor):

Dr. Jean Michel Maes (Editor General, RNB)
Museo Entomológico / Morpho Residency
De hielera CELSA media cuadra arriba
21000 León, NICARAGUA
Teléfono (505) 7791-2686
jmmaes@yahoo.com

También se puede remitir a los miembros del comité editorial de la revista.

Costos de publicación y sobretiros.

La publicación de un artículo es completamente gratis.

Los autores recibirán una versión PDF de su publicación para distribución.